

## What can I do to stay healthy?

- Let your doctor know if you have questions about your treatment or if you are not able to follow your prescribed treatment plan.
- Abstinence or mutual monogamy (having sex with only one partner who only has sex with you) is a way to avoid infection.
- Tell all of your sex partners that you have an STD, and recommend that they (and their other sex partners) be tested and treated. Your sex partner may not know they have a STD. However, neither you nor your partner need to have symptoms to be contagious.
- Until you and your sex partner(s) have successfully completed treatment, you can pass the infection to others during sexual contact even if you don't have any symptoms.
- Report any new symptoms to your doctor, particularly if you have any new pain or notice a change in the amount, appearance, or smell of your vaginal discharge. Many problems that affect a woman's reproductive tract, including sexually transmitted infections, can cause similar symptoms. It is important that you let your doctor evaluate your symptoms (even if you have had the same or similar symptoms before) so they can determine their exact cause, prescribe the right treatment, and teach you how to take steps to keep from spreading an infection to others.
- Protect your sex partners from becoming infected with a STD. Use latex condoms whenever you engage in sexual activity. Condoms may reduce your risk of passing the disease to your sex partner, but they will not protect you or your partner against all STDs. Only the skin that is covered by or comes in contact with the condom is protected from infection. Any uncovered skin on the genitals, groin, thighs, anus and rectum, and possibly in the mouth, is prone to infection.
- Visit your doctor as recommended for regular pelvic exams and screening tests. Keep a record of the dates and results of such exams and tests; it might come in handy if you ever change doctors or insurance providers or if you experience a reproductive health problem in the future.

### Where can I find out more about chlamydia and gonorrhea?

You can obtain additional information about these STDs from the following sources:

**Centers for Disease Control and Prevention** - Division of STD Prevention  
National STD hotline: (800) 232-4636 | [cdc.gov/std](http://cdc.gov/std)

**American Sexual Health Association (ASHA)**  
Telephone: (800) 783-9877 | [ashastd.org](http://ashastd.org)

#### References

1. Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines, 2015. *MMWR*. 2015;64(RR3):1-138.
2. American Sexual Health Association. Chlamydia. Available at: <http://www.ashsexualhealth.org/stdsstis/chlamydia/>. Accessed July 15, 2015.
3. Centers for Disease Control and Prevention. Chlamydia CDC fact sheet. Available at: <http://www.cdc.gov/std/chlamydia/stdfact-chlamydia.htm>. Accessed July 15, 2015.
4. American Sexual Health Association. Gonorrhea. Available at: <http://www.ashsexualhealth.org/stdsstis/gonorrhea/>. Accessed July 15, 2015.
5. Centers for Disease Control and Prevention. Gonorrhea CDC fact sheet. Available at: <http://www.cdc.gov/std/gonorrhea/stdfact-gonorrhea.htm>. Accessed July 15, 2015.

**Note:** This material is provided for general information purposes only. It is not intended as a substitute for medical advice and/or consultation with a physician or technical expert.



## PATIENT INFORMATION

# Chlamydia and Gonorrhea





## What are chlamydia and gonorrhea?

Chlamydia and gonorrhea are all common sexually transmitted diseases (STDs) that can cause infections in the genitals, rectum and throat. These diseases are easily spread by having vaginal, anal or oral sex with someone who is infected with the disease.

### Chlamydia

Chlamydia is a disease caused by infection with the bacteria called *Chlamydia trachomatis*.<sup>1,2</sup>

You may never develop symptoms of a chlamydial infection.<sup>1-3</sup> When chlamydia symptoms do occur, they are usually mild and appear 1 to 3 weeks after you become infected with *Chlamydia trachomatis*.<sup>1-3</sup> Symptoms include an abnormal vaginal discharge and pain when you urinate. If your male sex partner has chlamydia, he may experience an abnormal discharge from his penis and pain or burning during urination.

### Testing for chlamydia

Because symptoms of chlamydia are usually mild or may not occur at all, you should be tested annually for the infection if you are a sexually active woman age 25 and under.<sup>1-3</sup> Yearly testing is also recommended for women over age 25 that have risk factors for chlamydia (e.g., those with new partners and those with multiple sex partners, a sex partner with concurrent partners or a sex partner who has an STD).<sup>1-3</sup>

### Potential complications of chlamydia

If the infection is not diagnosed and treated early, chlamydia can result in complications like pelvic inflammatory disease (PID), infertility, ectopic pregnancy and more.

### Gonorrhea and its symptoms

Gonorrhea is a common disease that is passed from one person to another during sexual contact. It is caused by infection with bacteria called *Neisseria gonorrhoeae*.<sup>1,4</sup>

While many men exhibit symptoms of a gonorrheal infection, women usually do not.<sup>1,4-5</sup> When gonorrhea symptoms do occur, they include pain or burning during urination and a yellowish discharge.<sup>1,4-5</sup>

### Potential complications of gonorrhea

If a gonorrheal infection is not promptly treated, complications may include PID, ectopic pregnancies, miscarriage, infections in your newborn baby and more.

### Testing for gonorrhea

If you are younger than 25 years of age and are sexually active, you should receive an annual screening test for gonorrhea.<sup>1,4,5</sup> If you are older and have risk factors (e.g., those who have a new sex partner, more than one sex partner, a sex partner with concurrent partners or a sex partner who has an STD), you too should be screened.<sup>1,4,5</sup>

### How are chlamydia and gonorrhea diagnosed?

Sexually transmitted diseases are diagnosed using laboratory tests. Your healthcare provider may ask you to provide a urine sample or may use a swab to get a sample from your vagina or endocervical canal (the narrow passageway that leads from the cervix to the uterus) for testing. If you have symptoms that may indicate an infection in your throat and/or rectum, your doctor will obtain a sample of cells from those areas by rubbing them with a swab.

These samples are sent to Labcorp where tests will be performed to determine whether a STD is present. If you have symptoms of PID, your doctor may perform additional exams and tests to determine the extent of damage to your reproductive organs.

### How are chlamydia and gonorrhea treated?

These STDs are treated with antibiotics.<sup>1-5</sup> The drugs are usually taken by mouth but also may include an injection. The choice of drug and the length of treatment will depend on the severity of your infection, whether you are pregnant or breast-feeding or have ever had an allergic reaction to an antibiotic, and whether you have been treated for sexually transmitted infections in the past.

It is very important that you take all of your prescribed medication, even if your symptoms go away before you have taken all the pills. If you stop taking your medication after your symptoms are relieved but before you complete the full course of treatment, the infection may not be completely cured and may flare up again. It is also essential that all of your sex partners be tested and, if necessary, treated for infection. Medication should not be shared. Sexual contact should be avoided until you have been treated and cured. Follow your doctor's instructions on when you can have sexual contact again.

It is common for women who have been treated for STDs to develop another infection within several months. Most posttreatment infections are the result of reinfection caused by having sex with an infected partner. Because repeat infections may increase your risk for PID and other complications, your doctor is likely to recommend that you be retested for chlamydia and gonorrhea infection, preferably about 3 months after you have completed antibiotic treatment, but certainly within 12 months.<sup>1,2,3</sup>