

Labcorp Common ICD-10-CM Sign and Symptom Codes

Listed below are the ICD-10-CM codes commonly received by Labcorp from ordering physicians. Code descriptions were taken from the official code set, issued by the US Department of Health and Human Services, as updated effective October 1, 2024 through September 30, 2025. Depending on the nature of your practice, this list may be a useful reference tool; however, it is not a complete list. Please refer to an ICD-10-CM manual for a complete listing. Responsibility for the provision of correct diagnosis code(s) is with the ordering physician. The ordering physician must always determine, for the specific date of service, the appropriate diagnosis code(s) based on the patient's signs and symptoms. ICD-10-CM codes in effect at the time of service are required by third-party payers to support the medical necessity of the test(s) and/or profile(s) ordered. Physicians are asked to submit all medically appropriate ICD-10-CM codes when ordering laboratory testing.

Labcorp's provision of this information as a customer service does not suggest and is not meant to suggest that any of these codes should or should not be used on any given occasion. Labcorp makes no recommendation regarding the use of any particular diagnosis code(s). Diagnosis information should be reflected in the patient's medical record.

Symbol	Diagnosis	
	Code	Sign or Symptom
∩§	R79.9	Abnormal finding of blood chemistry, unspecified
	R74.8	Abnormal levels of other serum enzyme
§	R94.5	Abnormal results of liver function studies
§	R94.6	Abnormal results of thyroid function studies
§♀	N93.9	Abnormal uterine and vaginal bleeding, unspecified
§	R63.5	Abnormal weight gain
§	R63.4	Abnormal weight loss
§	L70.0	Acne vulgaris
➤§	N30.00	Acute cystitis without hematuria
★§	J02.9	Acute pharyngitis, unspecified
★←§	J06.9	Acute upper respiratory infection, unspecified
➤§♀	N76.0	Acute vaginitis
◀§	M81.0	Age-related osteoporosis without current pathological fracture
o	D63.1	Anemia in chronic kidney disease (never a primary diagnosis)
§	D64.9	Anemia, unspecified
	F41.9	Anxiety disorder, unspecified
‡♠	I25.10	Atherosclerotic heart disease of native coronary artery without angina pectoris
§♀	R87.610	Atypical squamous cells of undetermined significance on cytologic smear of smear (ASC-US)
§	E06.3	Autoimmune thyroiditis
◆§♂	N40.1	Benign prostatic hyperplasia with lower urinary tract symptoms
§♂	N40.0	Benign prostatic hyperplasia without lower urinary tract symptoms
§	N20.0	Calculus of kidney
§	R53.82	Chronic fatigue, unspecified
✦	N18.2	Chronic kidney disease, stage 2 (mild)
✦	N18.30	Chronic kidney disease, stage 3 unspecified
✦	N18.31	Chronic kidney disease, stage 3A
✦	N18.32	Chronic kidney disease, stage 3B
✦	N18.4	Chronic kidney disease, stage 4 (severe)
✦	N18.9	Chronic kidney disease, unspecified
★≈§	J44.9	Chronic obstructive pulmonary disease, unspecified
§	Z20.2	Contact with an(suspected) exposure to infections with a predominantly sexual mode of transmission
	R68.82	Decreased libido (adult age 15-124)
§	E53.8	Deficiency of other specified B group vitamins
§	R19.7	Diarrhea, unspecified

Symbol	Diagnosis	
	Code	Sign or Symptom
§	E07.9	Disorder of thyroid, unspecified
§	R42	Dizziness and giddiness
§	R30.0	Dysuria
	R03.0	Elevated blood-pressure reading, without diagnosis of hypertension
♂	R97.20	Elevated prostate specific antigen [PSA]
§	D72.829	Elevated white blood cell count, unspecified
	R74.01	Elevation of levels of liver transaminase levels
§♀	Z36.9	Encounter for antenatal screening, unspecified (maternity age 9-64)
∩;§♀	Z01.419	Encounter for gynecological examination (general) (routine) without abnormal findings
§	Z01.818	Encounter for other preprocedural examination
§♀	Z36.89	Encounter for other specified antenatal screening (maternity age 9-64)
§	Z01.89	Encounter for other specified special examinations
♀	Z32.01	Encounter for pregnancy test, result positive (maternity age 9-64)
§	Z01.812	Encounter for preprocedural laboratory examination
§	Z13.6	Encounter for screening for cardiovascular disorders
§	Z13.1	Encounter for screening for diabetes mellitus
§	Z13.0	Encounter for screening for disease of the blood and blood-forming organs and certain disorders involving the immune mechanism
§	Z11.4	Encounter for screening for human immunodeficiency virus [HIV]
§	Z11.51	Encounter for screening for human papillomavirus (HPV)
§	Z11.3	Encounter for screening for infections with a predominantly sexual mode of transmission
§	Z13.220	Encounter for screening for lipid disorders
♣§♀	Z12.4	Encounter for screening for malignant neoplasm of cervix
♣§	Z12.11	Encounter for screening for malignant neoplasm of colon
♣§♂	Z12.5	Encounter for screening for malignant neoplasm of prostate
§	Z13.21	Encounter for screening for nutritional disorder
§	Z13.89	Encounter for screening for other disorder
§	Z11.8	Encounter for screening for other infectious and parasitic diseases
§	Z13.228	Encounter for screening for other metabolic disorders
§	Z13.29	Encounter for screening for other suspected endocrine disorder
§	Z11.59	Encounter for screening for other viral diseases

Symbol	Diagnosis	
	Code	Sign or Symptom
§	Z11.1	Encounter for screening for respiratory tuberculosis
§	Z13.9	Encounter for screening, unspecified
§♀	Z34.9Ø	Encounter for supervision of normal pregnancy, unspecified, unspecified trimester (maternity age 9-64)
§♀	Z34.81	Encounter for supervision of other normal pregnancy, first trimester (maternity age 9-64)
§♀	Z34.82	Encounter for supervision of other normal pregnancy, secondary trimester (maternity age 9-64)
∞■§	Z51.81	Encounter for therapeutic drug level monitoring
▼§	E34.9	Endocrine disorder, unspecified
§	R1Ø.13	Epigastric pain
■§	I1Ø	Essential (primary) hypertension
§	K76.Ø	Fatty (change of) liver, not elsewhere classified
§	R5Ø.9	Fever, unspecified
↑§	R35.Ø	Frequency of micturition
§	K21.9	Gastro-esophageal reflux disease without esophagitis
	F41.1	Generalized anxiety disorder
★	M1Ø.9	Gout, unspecified
	R51.9	Headache, unspecified
§	R31.9	Hematuria, unspecified
§	Z72.51	High risk heterosexual behavior
*Ø§	B2Ø	Human immunodeficiency virus [HIV] disease
§	E83.52	Hypercalcemia
§	R73.9	Hyperglycemia, unspecified
§	E87.5	Hyperkalemia
§	E78.5	Hyperlipidemia, unspecified
■Δ§	I12.9	Hypertensive chronic kidney disease with stage 1 through stage 4 chronic kidney disease, or unspecified chronic kidney disease
§	E87.6	Hypokalemia
§	E87.1	Hypo-osmolality and hyponatremia
§	EØ3.9	Hypothyroidism, unspecified
§	R73.Ø1	Impaired fasting glucose
	G47.ØØ	Insomnia, unspecified
▲§	E61.1	Iron deficiency
§	D5Ø.Ø	Iron deficiency anemia secondary to blood loss (chronic)
	D5Ø.9	Iron deficiency anemia, unspecified
§♀	N92.6	Irregular menstruation, unspecified
⊕❖	Z79.Ø1	Long term (current) use of anticoagulants
⊕❖	Z79.4	Long term (current) use of insulin
⊕❖§	Z79.891	Long term (current) use of opiate analgesic
§	M54.5Ø	Low back pain, unspecified
◆§♂	C61	Malignant neoplasm of prostate
→§♀	N95.1	Menopausal and female climacteric states
§	E78.2	Mixed hyperlipidemia
▼§	E66.Ø1	Morbid (severe) obesity due to excess calories
§	D48.5	Neoplasm of uncertain behavior of skin
↑§	R35.1	Nocturia
▼§	E66.9	Obesity, unspecified
§	F11.2Ø	Opioid dependence, uncomplicated
§	R73.Ø9	Other abnormal glucose
§⊙	G89.29	Other chronic pain
§	R53.83	Other fatigue

Symbol	Diagnosis	
	Code	Sign or Symptom
	R68.89	Other general symptoms and signs
§	E78.49	Other hyperlipidemia
	D5Ø.8	Other iron deficiency anemias
⊕❖	Z79.899	Other long term (current) drug therapy
§	R53.81	Other malaise
↓§	R79.89	Other specified abnormal findings of blood chemistry
§	EØ3.8	Other specified hypothyroidism
§♀	N89.8	Other specified noninflammatory disorders of vagina
▼§	E66.3	Overweight
	M25.5Ø	Pain in unspecified joint
§	RØØ.2	Palpitations
	I48.Ø	Paroxysmal atrial fibrillation
§	R1Ø.2	Pelvic and perineal pain
	Z76.89	Persons encountering health services in other specified circumstances
§	E89.Ø	Postprocedural hypothyroidism
§	R73.Ø3	Prediabetes
§	R8Ø.9	Proteinuria, unspecified
§	E78.ØØ	Pure hypercholesterolemia, unspecified
§	E78.1	Pure hyperglyceridemia
§	R21	Rash and other nonspecific skin eruption
§	MØ5.79	Rheumatoid arthritis with rheumatoid factor of multiple sites without organ or systems involvement
§	RØ6.Ø2	Shortness of breath
▲§♂	E29.1	Testicular hypofunction
§	D69.6	Thrombocytopenia, unspecified
§	EØ5.9Ø	Thyrotoxicosis, unspecified without thyrotoxic crisis or storm
●▶§	E11.22	Type 2 diabetes mellitus with diabetic chronic kidney disease
●§	E11.42	Type 2 diabetes mellitus with diabetic polyneuropathy
●§	E11.65	Type 2 diabetes mellitus with hyperglycemia
●◀§	E11.69	Type 2 diabetes mellitus with other specified complication
●§	E11.8	Type 2 diabetes mellitus with unspecified complications
●§	E11.9	Type 2 diabetes mellitus without complications
§	R1Ø.9	Unspecified abdominal pain
↓§	R82.9Ø	Unspecified abnormal findings in urine
	I48.91	Unspecified atrial fibrillation
	R39.9	Unspecified symptoms and signs involving the genitourinary system
➤§	N39.Ø	Urinary tract infection, site not specified
§	D51.9	Vitamin B12 deficiency anemia, unspecified
§	E55.9	Vitamin D deficiency, unspecified

Legend for Symbols

- ↓ Use additional code to identify any retained foreign body, if applicable (Z18.-)
- § Excludes1 (mutually exclusive diagnosis code) notation applies (please refer to the ICD-10-CM manual)
- ♀ Female diagnosis only
- Use additional code to identify infectious agent (B95-B97)
- ☆ Use additional code, where applicable, to identify:
 - exposure to environmental tobacco smoke (Z77.22)
 - exposure to tobacco smoke in the perinatal period (P96.81)
 - history of tobacco dependence (Z87.891)
 - occupational exposure to environmental tobacco smoke (Z57.31)
 - tobacco dependence (F17.-)
 - tobacco use (Z72.0)
- ← Use additional code (B95-B97) to identify infectious agent, if known, such as:
 - respiratory syncytial virus (RSV) (B97.4)
- ↖ Use additional code to identify:
 - major osseous defect, if applicable (M89.7-)
 - personal history of (healed) osteoporosis fracture, if applicable (Z87.310)
- Code first underlying chronic kidney disease (CKD) (N18.-)
- ‡ Code also the presence of hypertension (I10-I1A)
- Use additional code to identify:
 - chronic total occlusion of coronary artery (I25.82)
 - exposure to environmental tobacco smoke (Z77.22)
 - history of tobacco dependence (Z87.891)
 - occupational exposure to environmental tobacco smoke (Z57.31)
 - tobacco dependence (F17.-)
 - tobacco use (Z72.0)
- ♠ Use additional code, if applicable, to identify:
 - coronary atherosclerosis due to calcified coronary lesion (I25.84)
 - coronary atherosclerosis due to lipid rich plaque (I25.83)
- ◇ Use additional code for associated symptoms, when specified:
 - incomplete bladder emptying (R39.14)
 - nocturia (R35.1)
 - straining on urination (R39.16)
 - urinary frequency (R35.0)
 - urinary hesitancy (R39.11)
 - urinary incontinence (N39.4-)
 - urinary obstruction (N13.8)
 - urinary retention (R33.8)
 - urinary urgency (R39.15)
 - weak urinary stream (R39.12)
- ♂ Male diagnosis only
- ✦ Code first any associated:
 - diabetic chronic kidney disease (E08.22, E09.22, E10.22, E11.22, E13.22)
 - hypertensive chronic kidney disease (I12.-, I13.-)Use additional code to identify kidney transplant status, if applicable (Z94.0)
- ≈ Code also type of asthma, if applicable (J45.-)
- ☼ Use additional code:
 - for screening for human papillomavirus, if applicable (Z11.51)
 - for screening vaginal pap smear, if applicable (Z12.72)
 - to identify acquired absence of uterus, if applicable (Z90.71-)
- ♣ Use additional code to identify any family history of malignant neoplasm (Z80.-)
- ∞ Code also condition requiring care
- ▣ Code also any long-term (current) drug therapy (Z79.-)

Legend for Symbols *(continued)*

- ▼ Code also the underlying disorder, such as:
 - primary neuroendocrine tumors (C7A.-)
 - secondary neuroendocrine tumors (C7B.-)
- Use additional code to identify:
 - exposure to environmental tobacco smoke (Z77.22)
 - history of tobacco dependence (Z87.891)
 - occupational exposure to environmental tobacco smoke (Z57.31)
 - tobacco dependence (F17.-)
 - tobacco use (Z72.Ø)
- ↑ Code first, if applicable, any causal condition, such as:
 - enlarged prostate (N4Ø.1)
- ★ Use additional code to identify:
 - autonomic neuropathy in diseases classified elsewhere (G99.Ø)
 - calculus of urinary tract in diseases classified elsewhere (N22)
 - cardiomyopathy in diseases classified elsewhere (I43)
 - disorders of external ear in diseases classified elsewhere (H61.1-, H62.8-)
 - disorders of iris and ciliary body in diseases classified elsewhere (H22)
 - glomerular disorders in diseases classified elsewhere (NØ8)
- * Use additional code to identify resistance to antimicrobial drugs (Z16.-)
- ◇ Code first Human immunodeficiency virus [HIV] disease complicating pregnancy, childbirth and the puerperium, if applicable (O98.7-)
 - Use additional code(s) to identify all manifestations of HIV infection
- △ Use additional code to identify the stage of chronic kidney disease (N18.1-N18.4, N18.9)
- ▲ Use additional code for adverse effect, if applicable, to identify drug (T36-T5Ø with fifth or sixth character 5)
- ⊗ Code also any follow-up examination (ZØ8-ZØ9)
- ❖ Code also any therapeutic drug level monitoring (Z51.81)
- ◆ Use additional code, if applicable, to identify:
 - hormone sensitivity status (Z19.1-Z19.2)
 - rising PSA following treatment for malignant neoplasm of prostate (R97.21)
- Use additional code for associated symptoms
- ▼ Code first obesity complicating pregnancy, childbirth and the puerperium, if applicable (O99.21-)
 - Use additional code to identify body mass index (BMI), if known, for adults (Z68.1-Z68.45) or pediatrics (Z68.5-)
- ⊙ Code also related psychological factors associated with pain (F45.42)
- Use additional code to identify control using:
 - insulin (Z79.4)
 - injectable non-insulin antidiabetic drugs (Z79.85)
 - oral antidiabetic drugs (Z79.84)
 - oral hypoglycemic drugs (Z79.84)
- ▶ Use additional code to identify stage of chronic kidney disease (N18.1-N18.6)
- ◀ Use additional code to identify complication